

Chapter 1 Motion in a Straight Line

Assignment 3

Class 11

Prerna Edu

PRERNA EDUCATION

DPP


DAILY PRACTICE PROBLEMS

CLASS : XITH
DATE :

SUBJECT : PHYSICS
DPP NO. : 3

Topic :- MOTION IN A STRAIGHT LINE


1. An object is dropped from rest. Its $v-t$ graph is



2. A particle is thrown vertically upwards. If its velocity at half of the maximum height is 10 m/sec , then maximum height attained by it is (Take $g = 10 \text{ m/sec}^2$)

a) 8 m b) 10 m c) 12 m d) 16 m

3. Which graph represents the uniform acceleration



4. What is the relation between displacement, time and acceleration in case of a body having uniform acceleration

a) $S = ut + \frac{1}{2}ft^2$ b) $S = (u + f) t$ c) $S = v^2 - 2fs$ d) None of these

5. The acceleration ' a ' in m/s^2 of a particle is given by $a = 3t^2 + 2t + 2$ where t is the time. If the particle starts out with a velocity $u = 2 \text{ m/s}$ at $t = 0$, then the velocity at the end of 2 seconds is

a) 12 m/s b) 18 m/s c) 27 m/s d) 36 m/s

PRERNA EDUCATION

6. Two bodies are thrown simultaneously from a tower with same initial velocity v_0 : one vertically upwards, the other vertically downwards. The distance between the two bodies after time t is
- a) $2v_0t + \frac{1}{2}gt^2$ b) $2v_0t$ c) $v_0t + \frac{1}{2}gt^2$ d) v_0t
7. An aeroplane flies 400 m north and 300 m south and then flies 1200 m upwards then net displacement is
- a) 1200 m b) 1300 m c) 1400 m d) 1500 m
8. The displacement of a particle undergoing rectilinear motion along the x -axis is given by $x = (2t^2 + 21t^2 + 60t + 6)$. The acceleration of the particle when its velocity is zero is
- a) 36ms^{-2} b) 9ms^{-2} c) -9ms^{-2} d) -18ms^{-2}
9. A river is flowing from W to E with a speed of 5 m/min. A man can swim in still water with a velocity 10 m/min. In which direction should the man swim so as to take the shortest possible path to go to the south
- a) 30° with downstream
b) 60° with downstream
c) 120° with downstream
d) South
10. The numerical ratio of displacement to the distance covered is always
- a) Less than one b) Equal to one
c) Equal to or less than one d) Equal to or greater than one
11. From the top of tower, a stone is thrown up. It reaches the ground in t_1 second. A second stone thrown down with the same speed reaches the ground in t_2 second. A third stone released from rest reaches the ground in t_3 second. Then

PRERNA EDUCATION

a) $t_3 = \frac{(t_1 + t_2)}{2}$ b) $t_3 = \sqrt{t_1 t_2}$ c) $\frac{1}{t_3} = \frac{1}{t_1} - \frac{1}{t_2}$ d) $t_3^2 = t_2^2 - t_1^2$

12. One car moving on a straight road covers one third of the distance with 20 km/hr and the rest with 60 km/hr. The average speed is

a) 40 km/hr b) 80 km/hr c) $46\frac{2}{3}$ km/hr d) 36 km/hr

13. A particle starts from rest, acceleration at 2 m/s^2 for 10 s and then goes with constant speed for 30 s and then decelerates at 4 m/s^2 till it stops. What is the distance travelled by it

a) 750 m b) 800 m c) 700 m d) 850 m

14. Acceleration of a particle changes when

- a) Direction of velocity changes b) Magnitude of velocity changes
c) Both of above d) Speed changes

15. A cat moves from X to Y with a uniform speed v_u and returns to X with a uniform speed v_d . The average speed for this ground trip is

a) $\frac{2v_d v_u}{v_d + v_u}$ b) $\sqrt{v_u v_d}$ c) $\frac{v_d v_u}{v_d + v_u}$ d) $\frac{v_u + v_d}{2}$

16. A boat takes two hours to travel 8 km and back in still water. If the velocity of water 4 kmh^{-1} , the time taken for going ups stream 8km and coming back is

- a) 2h b) 2 h 40 min
c) 1 h 20 min d) Cannot be estimated with the information given

17. A person travels along a straight road for the first half time with a velocity v_1 and the next half time with a velocity v_2

The mean velocity V of the man is

a) $\frac{2}{V} = \frac{1}{v_1} + \frac{1}{v_2}$ b) $V = \frac{v_1 + v_2}{2}$ c) $V = \sqrt{v_1 v_2}$ d) $V = \sqrt{\frac{v_1}{v_2}}$

18. A particle is projected with velocity v_0 along $x - axis$. The deceleration on the particle is proportional to the square of the distance from the origin i.e., $a = -ax^2$. The distance at which the particle stops is

PRERNA EDUCATION

a) $\sqrt{\frac{3v_0}{2\alpha}}$

b) $\left(\frac{3v_0}{2\alpha}\right)^{\frac{1}{3}}$

c) $\sqrt{\frac{3v_0^2}{2\alpha}}$

d) $\left(\frac{3v_0^2}{2\alpha}\right)^{\frac{1}{3}}$

19. Two balls are dropped to the ground from different heights. One ball is dropped 2 s after the other but they both strike the ground at the same time. If the first ball takes 5 s to reach the ground, then the difference in initial heights is ($g = 10 \text{ ms}^{-2}$)

a) 20 m

b) 80 m

c) 170 m

d) 40 m

20. A body starts from origin and moves along x -axis such that at any instant velocity is $v_t = 4t^3 - 2t$ where t is in second and v_t in ms^{-1} . The acceleration of the particle when it is 2m from the origin is

a) 28ms^{-2}

b) 22ms^{-2}

c) 12ms^{-2}

d) 10ms^{-2}