

CLASS : XIth
DATE :

SUBJECT : MATHS
DPP NO. :1

Topic :-SEQUENCES AND SERIES

1. If $p, q, r, s \in N$ and they are four consecutive terms of an A.P., then p th, q th, r th and s th terms of a G.P. are in

- a) A.P. b) G.P. c) H.P. d) None of these

2. $\frac{1 \cdot 2}{1^3} + \frac{2 \cdot 3}{1^3 + 2^3} + \frac{3 \cdot 4}{1^3 + 2^3 + 3^3} + \dots + n$ terms equals

- a) $\left(\frac{n}{n+1}\right)^2$ b) $\left(\frac{n}{n+1}\right)^3$ c) $\left(\frac{n}{n+1}\right)$ d) $\left(\frac{1}{n+1}\right)$

3. If $a_1, a_2, a_3, \dots, a_n$ are in AP, where $a_i > 0$ for all i , then value of $\frac{1}{\sqrt{a_1} + \sqrt{a_2}} + \frac{1}{\sqrt{a_2} + \sqrt{a_3}} + \dots + \frac{1}{a_{n-1} + \sqrt{a_n}}$ is equal to

- a) $\frac{n-1}{\sqrt{a_1} + \sqrt{a_n}}$ b) $\frac{n+1}{\sqrt{a_1} + \sqrt{a_n}}$ c) $\frac{n-1}{\sqrt{a_1} - \sqrt{a_n}}$ d) $\frac{n+1}{\sqrt{a_1} - \sqrt{a_n}}$

4. If $y = 2x^2 - 1$, then $\frac{1}{x^2} + \frac{1}{2x^4} + \frac{1}{3x^6} + \dots \infty$ equals to

- a) $\log_e \left(\frac{y+1}{y-1}\right)$ b) $\log_e \left(\frac{1+y}{1-y}\right)$ c) $\log_e \left(\frac{1-y}{1+y}\right)$ d) $\log \left(\frac{1+2y}{1-2y}\right)$

5. The interior angles of a polygon are in AP. If the smallest angle be 120° and the common difference be 5, then the number of side is

- a) 8 b) 10 c) 9 d) 6

6. If $\log_x(4x^{\log_5 x} + 5) = 2\log_5 x$, then x equals to

- a) 4, 5 b) -1, 5 c) 4, -1 d) $5, \frac{1}{5}$

7. Let a, b, c be in AP. If $0 < a, b, c < 1$, $x = \sum_{n=0}^{\infty} a^n$, $y = \sum_{n=0}^{\infty} b^n$ and $z = \sum_{n=0}^{\infty} c^n$, then

- a) $2y = x + z$ b) $2x = y + z$ c) $2z = x + y$ d) $2xz = xy + yz$

8. If $x^{\frac{3}{2}(\log_2 x - 3)} = \frac{1}{8}$, then x equals to

- a) 2 b) 3 c) 5 d) 6

9. If every terms of a GP with positive terms is the sum of its two previous terms, then the common ratio of the series is

- a) 1 b) $\frac{2}{\sqrt{5}}$ c) $\frac{\sqrt{5}-1}{2}$ d) $\frac{\sqrt{5}+1}{2}$

10. If $n_1, n_2, n_3, \dots, n_{100}$ are positive real numbers such that $n_1 + n_2 + n_3 + \dots + n_{100} = 20$ And $k = n_1(n_2 + n_3 + n_4)(n_5 + n_6 + \dots + n_9)(n_{10} + \dots + n_{16}) \dots (\dots + n_{100})$, then k belongs to
a) $(0, 100]$ b) $(0, 128]$ c) $[0, 144]$ d) None of these
11. If a, b, c are in AP, then the straight line $ax + by + c = 0$ will always pass through the point
a) $(-1, -2)$ b) $(1, -2)$ c) $(-1, 2)$ d) $(1, 2)$
12. If $\frac{e^x}{1-x} = B_0 + B_1x + B_2x^2 + \dots + B_nx^n + \dots$, then $B_n - B_{n-1}$ equals
a) $\frac{1}{n!}$ b) $\frac{1}{(n-1)!}$ c) $\frac{1}{n!} - \frac{1}{(n-1)!}$ d) 1
13. If $\frac{a+bx}{a-bx} = \frac{b+cx}{b-cx} = \frac{c+dx}{c-dx}$ ($x \neq 0$), then a, b, c, d are in
a) AP b) GP c) HP d) None of these
14. If $\sum_{r=1}^{\infty} \frac{1}{(2r-1)^2} = \frac{\pi^2}{8}$, then $\sum_{r=1}^{\infty} \frac{1}{r^2}$ is equal to
a) $\frac{\pi^2}{24}$ b) $\frac{\pi^2}{3}$ c) $\frac{\pi^2}{6}$ d) None of these
15. Jairam purchased a house in Rs 15000 and paid Rs 5000 at once. Rest money he promised to pay in annual installment of Rs 1000 with 10% per annum interest. How much money is to be paid by Jairam?
a) Rs 21555 b) Rs 20475 c) Rs 20500 d) Rs 20700
16. If a, b, c are in A.P., then $a + \frac{1}{bc}b + \frac{1}{ca}c + \frac{1}{ab}$ are in
a) A.P. b) G.P. c) H.P. d) None of these
17. The sum of the series $\frac{12}{2!} + \frac{28}{3!} + \frac{50}{4!} + \frac{78}{5!} + \dots$ is
a) e b) $3e$ c) $4e$ d) $5e$
18. The sum of the series $\frac{1}{2}x^2 + \frac{2}{3}x^3 + \frac{3}{4}x^4 + \frac{4}{5}x^5 + \dots$ is
a) $\frac{x}{1+x} + \log(1+x)$ b) $\frac{x}{1-x} + \log(1-x)$ c) $-\frac{x}{1+x} + \log(1+x)$ d) None of these
19. The sum of the infinite series $\left(\frac{1}{3}\right)^2 + \frac{1}{3}\left(\frac{1}{3}\right)^4 + \frac{1}{5}\left(\frac{1}{3}\right)^6 + \dots$ is
a) $\frac{1}{4}\log_e 2$ b) $\frac{1}{2}\log_e 2$ c) $\frac{1}{6}\log_e 2$ d) $\frac{1}{4}\log_e \frac{3}{2}$
20. Let T_r , be r th term of an AP whose first term is a and common difference is d . If for some positive integers $m, n, m \neq n$, $T_m = \frac{1}{n}$ and $T_n = \frac{1}{m}$, then $a - d$ equals
a) 0 b) 1 c) $\frac{1}{mn}$ d) $\frac{1}{m} + \frac{1}{n}$