

Topic :- STRAIGHT LINES

- A straight line through $P(1,2)$ is such that its intercept between the axes is bisected at P . Its equation is
a) $x + 2y = 5$ b) $x - y + 1 = 0$ c) $x + y - 3 = 0$ d) $2x + y - 4 = 0$
- The incentre of the triangle formed by the lines $x = 0, y = 0$ and $3x + 4y = 12$ is at
a) $(1/2, 1/2)$ b) $(1, 1)$ c) $(1, 1/2)$ d) $(1/2, 1)$
- A pair of perpendicular straight lines passes through the origin and also through the point of intersection of the curve $x^2 + y^2 = 4$ with $x + y = a$. The set containing the value of 'a' is
a) $\{-2, 2\}$ b) $\{-3, 3\}$ c) $\{-4, 4\}$ d) $\{-5, 5\}$
- If pairs straight lines $x^2 - 2pxy - y^2 = 0$ and $x^2 - 2qxy - y^2 = 0$ be such that each pair bisects the angle between the other pair, then
a) $pq = 1$ b) $pq = -1$ c) $pq = 2$ d) $pq = -2$
- In a rhombus $ABCD$ the diagonals AC and BD intersect at the point $(3,4)$. If the point A is $(1,2)$ the diagonal BD has the equation
a) $x - y - 1 = 0$ b) $x + y - 1 = 0$ c) $x - y + 1 = 0$ d) $x + y - 7 = 0$
- The gradient of one of the lines of $ax^2 + 2hxy + by^2 = 0$ is twice that of the other, then
a) $h^2 = ab$ b) $h = a + b$ c) $8h^2 = 9ab$ d) $9h^2 = 8ab$
- The family of lines making an angle 30° with the line $\sqrt{3}y = x + 1$ is
a) $x = \lambda$ (λ is parameter) b) $y = -\sqrt{3}x + \lambda$ (λ is parameter)
c) $y = \sqrt{3}x + \lambda$ d) None of the above
- If the slope of one of the lines represented by $ax^2 + 2hxy + by^2 = 0$ be the square of the other, then $\frac{a+h}{h} + \frac{8h^2}{ab}$ is
a) 3 b) 4 c) 5 d) 6
- The equation $y^2 - x^2 + 2x - 1 = 0$, represents
a) A pair of st. lines b) A circle c) A parabola d) An ellipse
- The vertices of a ΔOBC are $(0, 0)$, $B(-3, -1)$ and $C(-1, -3)$. The equation of a line parallel to BC and intersecting sides OB and OC whose distance from the origin is $1/2$, is
a) $x + y + \frac{1}{2} = 0$ b) $x + y - \frac{1}{2} = 0$ c) $x + y - \frac{1}{\sqrt{2}} = 0$ d) $x + y + \frac{1}{\sqrt{2}} = 0$
- The angle between the line joining the points $(1, -2)$, $(3, 2)$ and the line $x + 2y - 7 = 0$ is
a) π b) $\pi/2$ c) $\pi/3$ d) $\pi/6$
- The equation $y^2 - x^2 + 2x - 1 = 0$ represents
a) A hyperbola b) An ellipse
c) A pair of straight lines d) A rectangular hyperbola

13. The equation to the bisecting the join of $(3, -4)$ and $(5,2)$ and having its intercepts on the x -axis and the y -axis in the ratio $2 : 1$ is
 a) $x + y - 3 = 0$ b) $2x - y = 9$ c) $x + 2y = 2$ d) $2x + y = 7$
14. $A(-5,0)$ and $B(3,0)$ are two of the vertices of a triangle ABC . Its area is 20 square cms. The vertex C lies on the line $x - y = 2$. The coordinates of C are
 a) $(-7, -5)$ or $(3,5)$ b) $(-3, -5)$ or $(-5,7)$ c) $(7,5)$ or $(3,5)$ d) $(-3, -5)$ or $(7,5)$
15. The point of concurrence of the lines $ax + by + c = 0$ and a, b, c satisfy the relation $3a + 2b + 4c = 0$ is
 a) $(\frac{3}{2}, \frac{1}{4})$ b) $(\frac{3}{4}, \frac{1}{4})$ c) $(\frac{3}{4}, \frac{1}{2})$ d) $(\frac{3}{2}, \frac{1}{2})$
16. The angle between the straight line $x - y\sqrt{3} = 5$ and $\sqrt{3}x + y = 7$ is
 a) 90° b) 60° c) 75° d) 30°
17. The equation $y = \pm \sqrt{3}x, y = 1$ are the sides of
 a) An equilateral triangle b) A right angled triangle
 c) An isosceles triangle d) An obtuse triangle
18. A line passes through the point of intersection of the lines $3x + y + 1 = 0$ and $2x - y + 3 = 0$ and makes equal intercepts with axes. Then, equation of the line is
 a) $5x + 5y - 3 = 0$ b) $x + 5y - 3 = 0$ c) $5x - y - 3 = 0$ d) $5x + 5y + 3 = 0$
19. The equation of the straight line which passes through the point $(1, -2)$ and cuts off equal intercepts from the axes will be
 a) $x + y = 1$ b) $x - y = 1$ c) $x + y + 1 = 0$ d) $x - y - 2 = 0$
20. The orthocenter of a triangle formed by the lines $x + y = 1, 2x + 3y = 6$ and $4x - y + 4 = 0$ lies in the
 a) Ist quadrant b) IInd quadrant c) IIIrd quadrant d) IVth quadrant

