


Topic :- MOTION IN A PLANE

- The angle of projection at which the horizontal range and maximum height of projectile are equal is
 - 45°
 - $\theta = \tan^{-1}(0.25)$
 - $\theta = \tan^{-1} 4$ or $(\theta = 76^\circ)$
 - 60°
- A body slides down a frictionless track which ends in a circular loop of diameter D . Then the minimum height h of the body in terms of D so that it may just complete the loop, is
 - $h = \frac{5}{2}D$
 - $h = \frac{3}{2}D$
 - $h = \frac{5}{4}D$
 - $h = 2D$
- A force $\vec{F} = 2\hat{i} + 2\hat{j}$ N displaces a particle through $\vec{S} = 2\hat{i} + 2\hat{k}$ m in 16 s. The power developed by \vec{F} is
 - 0.25 J s^{-1}
 - 25 J s^{-1}
 - 225 J s^{-1}
 - 450 J s^{-1}
- A sphere of mass m is tied to end of a string of length l and rotated through the other end along a horizontal circular path with speed v . The work done in full horizontal circle is
 - 0
 - $\left(\frac{mv^2}{l}\right) \cdot 2\pi l$
 - $mg \cdot 2\pi$
 - $\left(\frac{mv^2}{l}\right) \cdot (l)$
- Two projectile are thrown with the same initial velocity at angles α and $(90^\circ - \alpha)$ with the horizontal. The maximum heights attained by them are h_1 and h_2 respectively. Then $\frac{h_1}{h_2}$ is equal
 - $\sin^2 \alpha$
 - $\cos^2 \alpha$
 - $\tan^2 \alpha$
 - 1
- A particle P is at the origin starts with velocity $\vec{v} = (2\hat{i} - 4\hat{j})\text{ms}^{-1}$ with constant acceleration $(3\hat{i} - 5\hat{j})\text{ms}^{-2}$. After travelling for 2 s, its distance from the origin is
 - 10 m
 - 10.2 m
 - 9.8 m
 - 11.7 m


7. A small sphere is hung by a string fixed to a wall. The sphere is pushed away from the wall by a stick. The force acting on the sphere are shown in figure. Which of the following statements is wrong?



- a) $P = W \tan \theta$ b) $\vec{T} + \vec{P} + \vec{W} = 0$ c) $T^2 = P^2 + W^2$ d) $T = P + W$

8. A particle moves in a circle of radius 30cm. Its linear speed is given by $v = 2t$, where t is in second and v in ms^{-1} . Find out its, radial and tangential acceleration at $t = 3\text{s}$, respectively,
 a) $220 \text{ ms}^{-2}, 50 \text{ ms}^{-2}$ b) $100 \text{ ms}^{-2}, 5 \text{ ms}^{-2}$ c) $120 \text{ ms}^{-2}, 2 \text{ ms}^{-2}$ d) $110 \text{ ms}^{-2}, 10 \text{ ms}^{-2}$

9. A small particle of mass m is projected at an angle θ with the x -axis with an initial velocity v_0 in the x - y plane as shown in the figure. At a time $t < \frac{v_0 \sin \theta}{g}$, the angular momentum of the particle is



- a) $-mgv_0 t^2 \cos \theta \hat{j}$ b) $mgv_0 t \cos \theta \hat{k}$ c) $-\frac{1}{2} mgv_0 t^2 \cos \theta \hat{k}$ d) $\frac{1}{2} mgv_0 t^2 \cos \theta \hat{i}$

10. A body is thrown upward from the earth surface with velocity 5 m/s and from a planet surface with velocity 3 m/s . Both follow the same path. What is the projectile acceleration due to gravity on the planet

- a) 2 m/s^2 b) 3.5 m/s^2 c) 4 m/s^2 d) 5 m/s^2

11. An unbanked curve has a radius of 60 m. The maximum speed at which the car make a turn is (Take $\mu = 0.75$)

- a) 7 ms^{-1} b) 14 ms^{-1} c) 21 ms^{-1} d) 2.1 ms^{-1}

12. A fly wheel rotates about a fixed axis and slows down from 300 rpm to 100 rpm in 2 min. Then its angular retardation in rad/min^2 is
- a) $\frac{100}{\pi}$ b) 100 c) 100π d) 200π
13. A particle is projected with a velocity 200 ms^{-1} at an angle of 60° . At the highest point, it explodes into three particles of equal masses. One goes vertically upwards with a velocity 100 ms^{-1} , the second particle goes vertically downwards. What is the velocity of third particle?
- a) 120 ms^{-1} making 60° angle with horizontal b) 200 ms^{-1} making 60° angle with horizontal
c) 300 ms^{-1} d) 200 ms^{-1}
14. A car is moving on a circular path and takes a turn. If R_1 and R_2 be the reactions on the inner and outer wheels respectively, then
- a) $R_1 = R_2$ b) $R_1 < R_2$ c) $R_1 > R_2$ d) $R_1 \geq R_2$
15. If the vector $\vec{A} = 2\hat{i} + 4\hat{j}$ and $\vec{B} = 5\hat{i} + p\hat{j}$ are parallel to each other, the magnitude of \vec{B} is
- a) $5\sqrt{5}$ b) 10 c) 15 d) $2\sqrt{5}$
16. A body is revolving with a uniform speed v in a circle of radius r . The tangential acceleration is
- a) $\frac{v}{r}$ b) $\frac{v^2}{r}$ c) Zero d) $\frac{v}{r^2}$
17. A bridge is in the form of a semi-circle of radius 40 m. The greatest speed with which a motor cycle can cross the bridge without leaving the ground at the highest point is ($g = 10 \text{ ms}^{-2}$) (frictional force is negligibly small)
- a) 40 ms^{-1} b) 20 ms^{-1} c) 30 ms^{-1} d) 15 ms^{-1}
18. A car is moving with high velocity when it has a turn. A force acts on it outwardly because of
- a) Centripetal force b) Centrifugal force c) Gravitational force d) All the above
19. If time of flight of a projectile is 10 seconds. Range is 500 meters. The maximum height attained by it will be
- a) 125 m b) 50 m c) 100 m d) 150 m
20. A stone is projected with a velocity $20\sqrt{2} \text{ ms}^{-1}$ at an angle of 45° to the horizontal. The average velocity of stone during its motion from starting point to its maximum height is ($g = 10 \text{ ms}^{-2}$)
- a) $5\sqrt{5} \text{ ms}^{-1}$ b) $10\sqrt{5} \text{ ms}^{-1}$ c) 20 ms^{-1} d) $20\sqrt{5} \text{ ms}^{-1}$