


Topic :- OSCILLATIONS

1. A small sphere carrying a charge q is hanging in between two parallel plates by a string of length L . Time period of pendulum is T_0 . When parallel plates are charged, the time period changes to T .

The ratio T/T_0 is equal to




- a) $\left(\frac{g + \frac{qE}{m}}{g}\right)^{1/2}$ b) $\left(\frac{g}{g + \frac{qE}{m}}\right)^{3/2}$ c) $\left(\frac{g}{g + \frac{qE}{m}}\right)^{1/2}$ d) None of these

2. The bob of a simple pendulum executes simple harmonic motion in water with a period t , while the period of oscillation of the bob is t_0 in air. Neglecting frictional force of water and given that the density of the bob is $(4/3 \times 1000 \text{ kg} - \text{m}^3)$. What relationship between t and t_0 is true?

- a) $t = t_0$ b) $t = t_0/2$ c) $t = 2t_0$ d) $t = 4t_0$

3. As a body performs S.H.M., its potential energy U . Varies with time as indicated in



4. Two simple pendulum of length 0.5 m and 20 m respectively are given small linear displacement in one direction at the same time. They will again be in the phase when the pendulum of shorter length has completed... oscillations.

- a) 5 b) 1 c) 2 d) 3

5. A simple harmonic oscillator has a period of 0.01 s and an amplitude of 0.2 m . The magnitude of the velocity in msec^{-1} at the centre of oscillation is
 a) 20π b) 100 c) 40π d) 100π
6. A body has a time period T_1 under the action of one force and T_2 under the action of another force, the square of the time period when both the forces are acting in the same direction is
 a) $T_1^2 T_2^2$ b) T_1^2 / T_2^2 c) $T_1^2 + T_2^2$ d) $T_1^2 T_2^2 / (T_1^2 + T_2^2)$
7. For a simple pendulum the graph between L and T will be
 a) Hyperbola b) Parabola c) A curved line d) A straight line
8. A mass of 4 kg suspended from a spring of force constant 800 Nm^{-1} executes simple harmonic oscillations. If the total energy of the oscillator is 4 J , the maximum acceleration (in ms^{-2}) of the mass is
 a) 5 b) 15 c) 45 d) 20
9. A spring of force constant k is cut into two pieces such that one piece is double the length of the other. Then the long piece will have a force constant of
 a) $(2/3)k$ b) $(3/2)k$ c) $3k$ d) $6k$
10. There is a body having mass m and performing S.H.M. with amplitude a . There is a restoring force $F = -Kx$, where x is the displacement. The total energy of body depends upon
 a) K, x b) K, a c) K, a, x d) K, a, v
11. If a body of mass 0.98 kg is made to oscillate on a spring of force constant 4.84 N/m , the angular frequency of the body is
 a) 1.22 rad/s b) 2.22 rad/s c) 3.22 rad/s d) 4.22 rad/s
12. The amplitude of vibration of a particle is given by $a_m = (a_0) / (a\omega^2 - b\omega + c)$; where a_0, a, b and c are positive. The condition for a single resonant frequency is
 a) $b^2 = 4ac$ b) $b^2 > 4ac$ c) $b^2 = 5ac$ d) $b^2 = 7ac$
13. The period of oscillation of a simple pendulum of constant length at earth surface is T . Its period inside a mine is
 a) Greater than T b) Less than T c) Equal to T d) Cannot be compared
14. In a simple pendulum, the period of oscillation T is related to length of the pendulum l as
 a) $\frac{l}{T} = \text{constant}$ b) $\frac{l^2}{T} = \text{constant}$ c) $\frac{l}{T^2} = \text{constant}$ d) $\frac{l^2}{T^2} = \text{constant}$

15. Starting from the origin a body oscillates simple harmonically with a period of 2 s. After what time will its kinetic energy be 75% of the total energy?

- a) $\frac{1}{6}$ s b) $\frac{1}{4}$ s c) $\frac{1}{3}$ s d) $\frac{1}{12}$ s

16. A mass m is suspended from a spring of length l and force constant K . The frequency of vibration of the mass is f_1 . The spring is cut into two equal parts and the same mass is suspended from one of the parts. The new frequency of vibration of mass is f_2 . Which of the following relations between the frequencies is correct

- a) $f_1 = \sqrt{2}f_2$ b) $f_1 = f_2$ c) $f_1 = 2f_2$ d) $f_2 = \sqrt{2}f_1$

17. How does time period of a pendulum vary with length?

- a) \sqrt{l} b) $\sqrt{\frac{l}{2}}$ c) $\frac{1}{\sqrt{l}}$ d) $2l$


18. A particle is vibrating in a simple harmonic motion with an amplitude of 4 cm. At what displacement from the equilibrium position, is its energy half potential and half kinetic

- a) 1 cm b) $\sqrt{2}$ cm c) 3 cm d) $2\sqrt{2}$ cm

19. A simple pendulum has a time period T_1 when on the earth's surface and T_2 when taken to a height $2R$ above the earth's surface where R is the radius of the earth. The value of (T_1/T_2) is

- a) 1/9 b) 1/3 c) $\sqrt{3}$ d) 3

20. A ball of mass (m) 0.5 kg is attached to the end of a string having length (L) 0.5 m. The ball is rotated on a horizontal circular path about vertical axis. The maximum tension that the string can bear is 324 N. The maximum possible value of angular velocity of ball (in radian/s) is



- a) 9 b) 18 c) 27 d) 36