

Topic :-Application of Derivatives

1. The set $\{x^3 - 12x : -3 \leq x \leq 3\}$ is equal to
a) $\{x : -16 \leq x \leq 16\}$ b) $\{x : -12 \leq x \leq 12\}$ c) $\{x : -9 \leq x \leq 9\}$ d) $\{x : 0 \leq x \leq 10\}$
2. If $xy = a^2$ and $S = b^2x + c^2y$ where a, b and c are constants, then the minimum value of S is
a) abc b) $\sqrt{a}bc$ c) $2abc$ d) None of these
3. Let $g(x) = f(x) + f'(1-x)$ and $f''(x) < 0, 0 \leq x \leq 1$. Then
a) $g(x)$ increases on $[1/2, 1]$ and decreases on $[0, 1/2]$
b) $g(x)$ decreases on $[0, 1]$
c) $g(x)$ increases on $[0, 1]$
d) $g(x)$ increases on $[0, 1/2]$ and decreases on $[1/2, 1]$
4. Select the correct statement from (a), (b), (c), (d) The function $f(x) = xe^{1-x}$
a) Strictly increasing in the interval $(\frac{1}{2}, 2)$ b) Increasing in the interval $(0, \infty)$
c) Decreases in the interval $(0, 2)$ d) Strictly decreasing in the interval $(1, \infty)$
5. If $\frac{a_0}{n+1} + \frac{a_1}{n} + \frac{a_2}{n-1} + \dots + \frac{a_{n-1}}{2} + a_n = 0$. Then the function $f(x) = a_0x^n + a_1x^{n-1} + a_2x^{n-2} + \dots + a_n$ has in $(0, 1)$
a) At least one zero b) At most one zero c) Only 3 zeros d) Only 2 zeros
6. A particle is moving along the curve $x = at^2 + bt + c$. If $ac = h^2$, then the particle would be moving with uniform
a) Rotation b) Velocity c) Acceleration d) Retardation

7. The approximate value of $(33)^{1/5}$ is
- a) 2.0125 b) 2.1 c) 2.01 d) None of these
8. At an instant the diagonal of a square is increasing at the rate of 0.2cm/sec and the area is increasing at the rate of $6\text{cm}^2/\text{sec}$. At that moment its side is
- a) $\frac{30}{\sqrt{2}}$ cm b) $30\sqrt{2}$ cm c) 30 cm d) 15 cm
9. A missile is fired from the ground level rises x metres vertically upwards in t seconds where $x = 100t - \frac{25}{2}t^2$. The maximum height reached is
- a) 200 m b) 125 m c) 160 m d) 190 m
10. The intercepts made by the tangent to the curve $y = \int_0^x |t|dt$, which is parallel to the line $y = 2x$, on y -axis are equal to
- a) 1, -1 b) -2, 2 c) 3 d) -3
11. The function $f(x) = \tan x - x$
- a) Always increases
b) Always decreases
c) Never decreases
d) Some times increases and some times decreases
12. The maximum value of xy subject to $x + y = 8$, is
- a) 8 b) 16 c) 20 d) 24
13. The tangent to the curve $y = 2x^2 - x + 1$ is parallel to the line $y = 3x + 9$ at the point
- a) (3, 9) b) (2, -1) c) (2, 1) d) (1, 2)
14. The point P of the curve $y^2 = 2x^3$ such that the tangent at P is perpendicular to the line $4x - 3y + 2 = 0$ is given by
- a) (2, 4) b) $(1, \sqrt{2})$ c) $(1/2, -1/2)$ d) $(1/8, -1/16)$
15. If the parametric equation of a curve given by $x = e^t \cos t$, $y = e^t \sin t$, then the tangent to the curve at the point $t = \pi/4$ makes with axis of x the angle
- a) 0 b) $\pi/4$ c) $\pi/3$ d) $\pi/2$

16. All points on the curve $y^2 = 4a\left(x + a \sin \frac{x}{a}\right)$ at which the tangents are parallel to the axis of x lie on a
- a) Circle b) Parabola c) Line d) None of these
17. The point of inflexion for the curve $y = x^{5/2}$ is
- a) (1, 1) b) (0, 0) c) (1, 0) d) (0, 1)
18. The minimum value of $2x + 3y$, when $xy = 6$, is
- a) 12 b) 9 c) 8 d) 6
19. If $f(x) = x^2 - 2x + 4$ on $[1, 5]$, then the value of a constant c such that $\frac{f(5) - f(1)}{5 - 1} = f'(c)$, is
- a) 0 b) 1 c) 2 d) 3
20. Let a, b be two distinct roots of a polynomial $f(x)$. Then there exists at least one root lying between a and b of the polynomial
- a) $f(x)$ b) $f'(x)$ c) $f''(x)$ d) None of these

