

## Topic :-DIFFERENTIAL EQUATIONS

1. The solution of the differential equation  $x \frac{dy}{dx} = 2y + x^3 e^x$ , where  $y = 0$  when  $x = 1$ , is  
 a)  $y = x^3(e^x - e)$       b)  $y = x^3(e - e^x)$       c)  $y = x^2(e^x - e)$       d)  $y = x^2(e - e^x)$
  
2. The solution of  $(1 + x^2) \frac{dy}{dx} + 2xy - 4x^2 = 0$  is  
 a)  $3x(1 + y^2) = 4y^3 + c$       b)  $3y(1 + x^2) = 4x^3 + c$   
 c)  $3x(1 - y^2) = 4y^3 + c$       d)  $3y(1 + y^2) = 4x^3 + c$
  
3. A normal is drawn at a  $P(x,y)$  of a curve. It meets the  $x$ -axis at Q. if  $PQ$  is of constant length  $k$ , then the differential equation describing such a curve is  
 a)  $y \frac{dy}{dx} = \pm \sqrt{k^2 - y^2}$       b)  $x \frac{dy}{dx} = \pm \sqrt{k^2 - x^2}$       c)  $y \frac{dy}{dx} = \pm \sqrt{y^2 - k^2}$       d)  $x \frac{dy}{dx} = \pm \sqrt{x^2 - k^2}$
  
4. The solution of the differential equation  $y_1 y_3 = 3y_2^2$  is  
 a)  $x = A_1 y^2 + A_2 y + A_3$       b)  $x = A_1 y + A_2$       c)  $x = A_1 y^2 + A_2 y$       d) None of these
  
5. If  $x = A \cos 4t + B \sin 4t$ , then  $\frac{d^2 x}{dt^2}$  is equal to  
 a)  $-16x$       b)  $16x$       c)  $x$       d)  $-x$
  
6. The order of the differential equation associated with the primitive  $y = c_1 + c_2 e^x + c_3 e^{-2x+c_4}$ , where  $c_1, c_2, c_3, c_4$  are arbitrary constants, is  
 a) 3      b) 4      c) 2      d) None of these
  
7. The differential equation of all parabolas whose axes are parallel to axis of  $x$ , is  
 a)  $\frac{d^3 y}{dx^3} = 0$       b)  $\frac{d^3 x}{dy^3} = 0$       c)  $\frac{d^2 y}{dx^2} + \frac{dy}{dx} = 0$       d)  $\frac{d^2 x}{dy^2} = 0$
  
8. The solution of the differential equation  $(x^2 - yx^2) \frac{dy}{dx} + y^2 + xy^2 = 0$  is  
 a)  $\log\left(\frac{x}{y}\right) = \frac{1}{x} + \frac{1}{y} + c$       b)  $\log\left(\frac{y}{x}\right) = \frac{1}{x} + \frac{1}{y} + c$   
 c)  $\log(xy) = \frac{1}{x} + \frac{1}{y} + c$       d)  $\log(xy) + \frac{1}{x} + \frac{1}{y} = c$

9. The solution of the differential equation  $x dy - y dx - \sqrt{x^2 - y^2} dx = 0$  is
- a)  $y - \sqrt{x^2 + y^2} = cx^2$       b)  $y + \sqrt{x^2 + y^2} = cx^2$   
c)  $y + \sqrt{x^2 + y^2} = cy^2$       d)  $x - \sqrt{x^2 + y^2} = cy^2$
10. Solution of  $\frac{dy}{dx} = \frac{x \log x^2 + x}{\sin y + y \cos y}$  is
- a)  $y \sin y = x^2 \log x + c$       b)  $y \sin y = x^2 + c$   
c)  $y \sin y = x^2 + \log x$       d)  $y \sin y = x \log x + c$
11. If integrating factor of  $x(1 - x^2)dy + (2x^2y - y - ax^3)dx = 0$  is  $e^{\int P dx}$ , then  $P$  is equal to
- a)  $\frac{2x^2 - ax^3}{x(1 - x^2)}$       b)  $2x^2 - 1$       c)  $\frac{2x^2 - 1}{ax^3}$       d)  $\frac{2x^2 - 1}{x(1 - x^2)}$
12. The solution of the differential equation  $\frac{dy}{dx} + \frac{y}{x} = x^2$ , is
- a)  $y = \frac{x^2}{4} + C x^{-2}$       b)  $y = x^{-1} + C x^{-3}$       c)  $y = \frac{x^3}{4} + C x^{-1}$       d)  $xy = x^2 + C$
13. The differential equation of all circles passing through the origin and having their centres on the  $x$ -axis is
- a)  $x^2 = y^2 + xy \frac{dy}{dx}$       b)  $x^2 = y^2 + 3xy \frac{dy}{dx}$       c)  $y^2 = x^2 + 2xy \frac{dy}{dx}$       d)  $y^2 = x^2 - 2xy \frac{dy}{dx}$
14. If  $y'' - 3y' + 2y = 0$  where  $y(0) = 1$ ,  $y'(0) = 0$ , then the value of  $y$  at  $x = \log 2$  is
- a) 1      b) -1      c) 2      d) 0
15. The differential equation of all straight lines touching the circle  $x^2 + y^2 = a^2$  is
- a)  $(y - \frac{dy}{dx})^2 = a^2 [1 + (\frac{dy}{dx})^2]$       b)  $(y - x \frac{dy}{dx})^2 = a^2 [1 + (\frac{dy}{dx})^2]$   
c)  $(y - x \frac{dy}{dx}) = a^2 [1 + \frac{dy}{dx}]$       d)  $(y - \frac{dy}{dx}) = a^2 [1 - \frac{dy}{dx}]$
16. The solution of the differential equation  $(x^2 - yx^2) \frac{dy}{dx} + y^2 + xy^2 = 0$  is
- a)  $\log(\frac{x}{y}) = \frac{1}{x} + \frac{1}{y} + c$       b)  $\log(\frac{y}{x}) = \frac{1}{x} + \frac{1}{y} + c$       c)  $\log(xy) = \frac{1}{x} + \frac{1}{y} + c$       d)  $\log(xy) + \frac{1}{x} + \frac{1}{y} = c$
17. The equation of the curve satisfying the equation  $(xy - x^2) \frac{dy}{dx} = y^2$  and passing through the point  $(-1, 1)$  is
- a)  $y = (\log y - 1)x$       b)  $y = (\log y + 1)x$       c)  $x = (\log x - 1)y$       d)  $x = (\log x + 1)y$
18.  $y = 2e^{2x} - e^{-x}$  is a solution of the differential equation
- a)  $y_2 + y_1 + 2y = 0$       b)  $y_2 - y_1 + 2y = 0$       c)  $y_2 + y_1 = 0$       d)  $y_2 - y_1 - 2y = 0$

19. The solution of  $y' - y = 1$ ,  $y(0) = -1$  is given by  $y(x)$ , which is equal to  
a)  $-\exp(x)$                       b)  $-\exp(-x)$                       c)  $-1$                       d)  $\exp(x) - 2$
20. The differential equation of the family of circles with fixed radius 5 unit and centre on the line  $y = 2$ , is  
a)  $(x - 2)^2 y'^2 = 25 - (y - 2)^2$                       b)  $(x - 2) y'^2 = 25 - (y - 2)^2$   
c)  $(y - 2) y'^2 = 25 - (y - 2)^2$                       d)  $(y - 2)^2 y'^2 = 25 - (y - 2)^2$

PE