

CLASS : XIIth  
DATE :

SUBJECT : MATHS  
DPP NO. : 2

**Topic :-INVERSE TRIGONOMETRIC FUNCTIONS**

1. If  $x_1, x_2, x_3, x_4$  are the roots of the equation  $x^4 - x^3 \sin 2\beta - x \cos \beta - \sin \beta = 0$ , then  $\tan^{-1} + \tan^{-1} x_2 + \tan^{-1} x_3 + \tan^{-1} x_4$  is equal to  
 a)  $\beta$                                   b)  $\frac{\pi}{2} - \beta$                                   c)  $\pi - \beta$                                   d)  $-\beta$
2. If  $x \in (-\frac{\pi}{2}, \frac{\pi}{2})$ , then the value of  $\tan^{-1}\left(\frac{\tan x}{4}\right) + \tan^{-1}\left(\frac{3 \sin 2x}{5 + 3 \cos 2x}\right)$  is  
 a)  $\frac{x}{2}$                                     b)  $2x$     c)  $3x$     d)  $x$
3.  $\frac{\alpha^3}{2} \operatorname{cosec}^2\left(\frac{1}{2} \tan^{-1} \frac{\alpha}{\beta}\right) + \frac{\beta^3}{2} \sec^2\left(\frac{1}{2} \tan^{-1} \left(\frac{\beta}{\alpha}\right)\right)$  is  
 a)  $(\alpha - \beta)(\alpha^2 + \beta^2)$     b)  $(\alpha + \beta)(\alpha^2 - \beta^2)$     c)  $(\alpha + \beta)(\alpha^2 + \beta^2)$     d) None of these
4. If  $-1 \leq x \leq 0$ , then  $\cos^{-1}(2x^2 - 1)$  equals  
 a)  $2\cos^{-1}x$                               b)  $\pi - 2\cos^{-1}x$                               c)  $2\pi - 2\cos^{-1}x$                               d)  $-2\cos^{-1}x$
5. If  $\cos^{-1}\frac{3}{5} - \sin^{-1}\frac{4}{5} = \cos^{-1}x$ , then  $x$  is equal to  
 a) 0    b) 1    c)  $-1$     d) None of these
6. If  $\sec^{-1}x = \operatorname{cosec}^{-1}y$ , then  $\cos^{-1}\frac{1}{x} + \cos^{-1}\frac{1}{y} =$   
 a)  $\pi$     b)  $\frac{\pi}{4}$     c)  $-\frac{\pi}{2}$     d)  $\frac{\pi}{2}$
7.  $\sec^2(\tan^{-1} 2) + \operatorname{cosec}^2(\cot^{-1} 3)$  is equal to  
 a) 1    b) 5    c) 10    d) 15
8. If  $-1 \leq x \leq -\frac{1}{2}$ , then  $\sin^{-1}(3x - 4x^3)$  equals  
 a)  $3\sin^{-1}x$                                 b)  $\pi - 3\sin^{-1}x$                                 c)  $-\pi - 3\sin^{-1}x$                             d) None of these
9.  $\tan\frac{2\pi}{5} - \tan\frac{\pi}{15} - \sqrt{3}\tan\frac{2\pi}{5}\tan\frac{\pi}{15}$  is equal to  
 a)  $-\sqrt{3}$                                     b)  $\frac{1}{\sqrt{3}}$     c) 1    d)  $\sqrt{3}$

10. The value of  $\tan\left\{\cos^{-1}\left(-\frac{2}{7}\right) - \frac{\pi}{2}\right\}$  is

a)  $\frac{2}{3\sqrt{5}}$

b)  $\frac{2}{3}$

c)  $\frac{1}{\sqrt{5}}$

d)  $\frac{4}{\sqrt{5}}$

11. The value of

$$\sin\left(\sin^{-1}\frac{1}{3} + \sec^{-1} 3\right) + \cos\left(\tan^{-1}\frac{1}{2} + \tan^{-1} 2\right)$$
 is

a) 1

b) 2

c) 3

d) 4

12. If  $-\frac{1}{\sqrt{3}} < x < \frac{1}{\sqrt{3}}$ , then  $\tan^{-1}\left(\frac{3x-x^3}{1-3x^2}\right)$  equals

a)  $3\tan^{-1}x$

b)  $-\pi + 3\tan^{-1}x$

c)  $\pi + 3\tan^{-1}x$

d) None of these

13.  $\sin\left(\frac{1}{2}\cos^{-1}\frac{4}{5}\right) =$

a)  $-\frac{1}{\sqrt{10}}$

b)  $\frac{1}{\sqrt{10}}$

c)  $-\frac{1}{10}$

d)  $\frac{1}{10}$

14. The solution of  $\tan^{-1} 2x + \tan^{-1} 3x = \frac{\pi}{4}$  is

a)  $\frac{1}{6}$

b) -1

c)  $(\frac{1}{6}, -1)$

d) None of these

15.  $\sin^{-1}\frac{4}{5} + 2\tan^{-1}\frac{1}{3}$  is equal to

a)  $\frac{\pi}{3}$

b)  $\frac{\pi}{4}$

c)  $\frac{\pi}{2}$

d) 0

16. The equation  $2\cos^{-1}x + \sin^{-1}x = \frac{11\pi}{6}$  has

a) No solution

b) Only one solution

c) Two solutions

d) Three solutions

17. The value of  $\cos^{-1}\left(\cos\frac{5\pi}{3}\right) + \sin^{-1}\left(\cos\frac{5\pi}{3}\right)$  is

a)  $\frac{10\pi}{3}$

b) 0

c)  $\frac{\pi}{2}$

d)  $\frac{5\pi}{3}$

18. The value of  $\sin^{-1}\left(\frac{\sqrt{3}}{2}\right) - \sin^{-1}\left(\frac{1}{2}\right)$  is

a)  $45^\circ$

b)  $90^\circ$

c)  $15^\circ$

d)  $30^\circ$

19. If  $\sin^{-1}x + \sin^{-1}(1-x) = \cos^{-1}x$ , then  $x$  equals

a) 1, -1

b) 1, 0

c)  $0, \frac{1}{2}$

d) None of these

20.  $\tan\left(\frac{\pi}{4} + \frac{1}{2}\cos^{-1}x\right) + \tan\left(\frac{\pi}{4} - \frac{1}{2}\cos^{-1}x\right), x \neq 0$  is equal to

a)  $x$

b)  $2x$

c)  $\frac{2}{x}$

d) None of these