

CLASS: XIIth DATE:

SUBJECT: MATHS

DPP NO.: 1

- 1. Let $A = \{x : -1 \le x \le 1\}$ and $f : A \to A$ such that f(x) = x|x|, then f is
 - a) A bijection
 - b) Injective but not surjective
 - c) Surjective but not injective
 - d) Neither injective nor surjective
- The domain of the function $\sin^{-}(\log_2 \frac{x^2}{2})$ is

3. If f(x) = ax + b and g(x) = cx + d, then f(g(x)) = g(f(x)) is equivalent to

a)
$$f(a) = f(c)$$

$$b) f(b) = g(b)$$
 $c) f(d) = g(b)$

c)
$$f(d) = g(b)$$

$$d) f(c) = g(a)$$

4. The period of the function $f(x) = \sin^4 3x + \cos^4 3x$ is

a)
$$\pi/2$$

b)
$$\pi/3$$

c)
$$\pi/6$$

d) None of these

5. Given $f(x) = \log_{10}\left(\frac{1+x}{1-x}\right)$ and $g(x) = \frac{3x+x^3}{1+3x^2}$, then $f \circ g(x)$ equals

a)
$$-f(x)$$

b)
$$3 f(x)$$

c)
$$[f(x)]^3$$

d) None of these

6. Which of the following functions is not an are not an injective map(s)?

a)
$$f(x) = |x + 1|, x \in [-1, \infty)$$

b)
$$g(x) = x + \frac{1}{x}$$
, $x ∈ (0, ∞)$

c)
$$h(x) = x^2 + 4x - 5, x \in (0, \infty)$$

$$d) h(x) = e^{-x}, x \in [0, \infty)$$

- 7. If $f:R \to R$ and $g:R \to R$ are defined by f(x) = x [x] and g(x) = [x] for $x \in R$, where [x] is the greatest integer not exceeding x, then for every $x \in R$, f(g(x)) is equal to
 - a) x

b)0

- c) f(x)
- d)g(x)

- The domain of definition of $f(x) = \sqrt{\frac{\log_{0.3}|x-2|}{|x|}}$, is
 - a) $[1, 2) \cup (2, 3]$
- b) [1, 3]
- c) R (1, 3]
- d) None of these

- 9. $f:R \rightarrow R$ given by $f(x) = 5 3\sin x$, is
 - a) One-one
- b) Onto
- c) One-one and onto
- d) None of these

10.	If $f(x + 2y, x - 2y) = x$ a) $\frac{x^2 - y^2}{8}$		v) equals	c) $\frac{x^2+y^2}{4}$		$d)\frac{x^2-y^2}{2}$
11.	If $f:R \to R$ is defined as $f(1-x)^{-1/3}$	$f(x) = (1 - x)^3$ b) $(1 - x)^3$	$(x)^{1/3}$, then f	$x^{-1}(x)$ is c) $1 - x^3$		d) $1 - x^{1/3}$
12.	If $f(x + 2y,x,x - 2y) =$ $a) \frac{x^2 - y^2}{8}$		x, y) equals	$c)\frac{x^2+y^2}{4}$		$\mathrm{d})\frac{x^2-y^2}{2}$
13.	Let $f:[4, \infty[\to [4, \infty[$ be defined by $f(x) = 5^{x(x-4)}]$ a) $2 - \sqrt{4 + \log_5 x}$ b) $2 + \sqrt{4 + \log_5 x}$			then $f^{-1}(x)$ c) $\left(\frac{1}{5}\right)^{x(x-4)}$		d) Not defined
14.	If $f:[2, 3] \to R$ is defined by $f(x) = x^3 + 3x - 2$, the a) [1, 12] b) [12, 34]			nen the range c) [35, 50]	f(x) is con	ntained in the interval d)[-12, 12]
15.	The period of $\sin^2 \theta$, is a) π^2	b) π		c) 2 π		$d)\pi/2$
16.	If $n \in N$, and the period of $\frac{\cos nx}{\sin(\frac{x}{n})}$ is 4π , then n is equal to					
	a) 4	b) 3		c) 2		d)1
17.	Foe real x , let $f(x) = x^3 + 5x + 1$, then a) f is one-one but not onto R c) f is one-one and onto R			b) <i>f</i> is onto <i>R</i> but not one-one d) <i>f</i> is neither one-one nor onto <i>R</i>		
18.	The range of the function	on $f(x) = \frac{1}{2}$	$\frac{1}{-\cos 3x}$, is			
	a) [- 1/3, 0]	b) <i>R</i>		c) [1/3, 1]		d) None of these
19. Let $A = \{2, 3, 4, 5,, 16, 17, 18\}$. Let be the equivalence relation on $A \times A$, cartesian product of A and A , defined by $(a, b) \approx (c, d)$ if $ad = bc$, then the number of ordered pairs of the equivalence class of $(3, 2)$ is						

a) 4

b) 5

c) 6

d)7

20. Let n be the natural number. Then, the range of the function $f(n) = 8 - n_{P_n-4}$, $4 \le n \le 6$, is

- a) {1, 2, 3, 4}
- b) {1, 2, 3, 4, 5, 6}
- c) {1, 2, 3}
- d){1, 2, 3, 4, 5}