

CBSE Test Paper 05 Chapter 10 Circle


1. If two tangents inclined at an angle of 60° are drawn to a circle of radius 3 cm, then the length of each tangent is equal to: (1)



- a. 6 cm
- b. $3\sqrt{3}$
- c. 3 cm
- d. $\frac{3}{2}\sqrt{3}$ cm
- 2. In the given fig., if O is the center of a circle, PQ is a chord and the tangent PR at P makes an angle of 50° with PQ, then \angle POQ is equal to : (1)




- a. 75°
- b. 100°
- c. 90°
- d. 80°
- **3.** In the given figure, O is the centre of the circle and PT is a tangent at T. If PC = 3 cm and PT = 6 cm, then the radius of the circle is equal to: **(1)**



a. 6 cm


- b. 5 cm
- c. 7 cm
- d. 4.5 cm

4. In the given figure, PQL and PRM are tangents to the circle with centre 0 at the points Q and R respectively and S is a point on the circle such that $\angle SQL = 50^o$ and $\angle SRM = 60^o$. Then $\angle QSR$ is equal to: (1)




- a. 50°
- b. 40°
- c. 60°
- d. 70°

5. In the given figure, if \triangle ABC is circumscribing a circle, then the length of BC is: (1)




- a. 10 cm
- b. 7 cm
- c. 11 cm
- d. 18 cm




- 7. If a circle can be inscribed in a parallelogram how will the parallelogram change? (1)
- **8.** What do you say about the line which is perpendicular to the radius of the circle through the point of contact? **(1)**
- **9.** How many common tangents can be drawn to two circles intersecting at two distinct points? **(1)**
- 10. In fig., PA and PB are two tangents drawn from an external point P to a circle with

centre C and radius 4 cm. If PA \perp PB, then find the length of each tangent. (1)




11. A tangent PQ at a point P on a circle of radius 5 cm meets a line through the centre 0 at a point Q so that OQ = 12 cm. Find the length PQ. **(2)**


12. In the given figure, two circles touch each other at the point C. Prove that the common tangent to the circles at C, bisects the common tangent at P and Q. **(2)**



13. In the given figure, 0 is the centre of a circle, BOA is its diameter and the tangent at the point P meets BA extended at T. If $\angle PBO = 30^{\circ}$, then find $\angle PTA$. (2)



14. In the given figure, 0 is the centre of a circle. PT and PQ are tangents to the circle from an external point P. If $\angle TPQ = 70^\circ$, find $\angle TRQ$. (3)



15. Prove that the tangent drawn at the midpoint of an arc of a circle is parallel to the chord joining the end points of the arc. **(3)**

16. Two concentric circles are of radii 5 cm and 3 cm. Find the length of the chord of the larger circle which touches the smaller circle. **(3)**

- **17.** A circle touches the sides of a quadrilateral ABCD at P, Q, R, S respectively. Show that the angles subtended at the centre by a pair of opposite sides are supplementary. **(3)**
- **18.** A triangle ABC is drawn to circumscribe a circle of radius 4 cm such that the segments BD and DC into which BC is divided by the point of contact D are of lengths 8 cm and 6 cm respectively. Find the sides AB and AC. **(4)**
- **19.** Prove that opposite sides of a quadrilateral circumscribing a circle subtend supplementary angles at the centre of the circle. **(4)**
- **20.** Two circles with centres O and O' of radii 3 cm and 4 cm, respectively intersect at two points P and Q such that OP and O'P are tangents to the two circles. Find the length of the common chord PQ. **(4)**

CBSE Test Paper 05 Chapter 10 Circle

Solution

1. b. $3\sqrt{3}$

Explanation: Refer fig

PQ and PR are two tangents to a circle

$$PQ = PR$$

PO bisects the angle between two tangents

therefore angle \angle OPQ = \angle OPR = 30°

In right angled triangle OPQ

$$\tan 30^\circ = \frac{OQ}{PQ}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{3}{PQ}$$

$$\Rightarrow$$
PQ = $3\sqrt{3}$ cm = PR

2. b. 100°

Explanation: Since OP is perpendicular to PR,

then
$$\angle \mathrm{OPR} = 90^{\circ}$$

$$\Rightarrow \angle RPQ + \angle QPO = 90^{\circ}$$

$$\Rightarrow 50^{\circ} + \angle QPO = 90^{\circ}$$

$$\Rightarrow \angle QPO = 40^{\circ}$$

Now, OP = OQ {Radii of same circle]

$$\therefore \angle OPQ = \angle OQP = 40^{\circ}$$
 [Angles opposite to equal sides]

In triangle OPQ,

$$\angle POQ + \angle OPQ + \angle OQP = 180^{\circ}$$

$$\Rightarrow$$
 \angle POQ + 40° + 40° = 180°

$$\Rightarrow$$
 \angle POQ = 100°

3. d. 4.5 cm

Explanation: In right angled triangle OTP,

Let the radius of the circle be r cm, then $\mathsf{OT} = \mathsf{OC} = \mathsf{r}$

$$OP^2 = OT^2 + PT^2$$

$$\Rightarrow (r+3)^2 = r^2 + 6^2$$
$$\Rightarrow r^2 + 6r + 9 = r^{2+36}$$
$$\Rightarrow 6r = 27 \Rightarrow r = 4.5cm$$

4. d. 70°

Explanation: Here
$$\angle OQS = \angle OQL - \angle SQL = 90^\circ - 50^\circ = 40^\circ$$

And $\angle ORS = \angle ORM - \angle SQM 90^\circ - 60^\circ = 30^\circ$ Since $OS = OQ$ [Radii] $\Rightarrow \angle OSQ = \angle OQS = 40^\circ$ [Angles opposite to equal sides] Again, since $OS = OR$ [Radii] $\Rightarrow \angle OSR = \angle ORS = 30^\circ$ [Angles opposite to equal sides] $\therefore \angle OSR = \angle OSQ + \angle OSR = 40^\circ + 30^\circ = 70^\circ$

5. a. 10 cm

Explanation: Let point of contact of tangent AB be P, point of contact of tangent BC be Q and point of contact of tangent AC be R.

Since, Tangents from an external points are equal.

$$\therefore BP = BQ = 3cm$$

$$PA = AR = 4 cm$$

$$\Rightarrow CR = 11 - 4 = 7 cm$$

$$CR = QC = 7 cm$$

$$\therefore BC = CQ + BQ = 7 + 3 = 10 cm$$


6. Circle touches both the parallel lines

Given, Distance between the parallel lines = 14 cm


We know that, Diameter of circle = Distance between the parallel lines

∴Radius =
$$\frac{14}{2}$$
 = 7 cm

- 7. It changes into a rectangle or a square.
- 8. The line which is perpendicular to the radius of the circle through the point of contact will be tangent to the circle. A line which intersects a circle at any one point is called the tangent.
- 9. 2 common tangents can be drawn to two circles intersecting at two distinct points.



10.



Construction: Join AC and BC

Now, $AC \perp AP$ and $CB \perp$

$$BP = 90^{0}$$


$$\angle APB =$$

Therefore, CAPB will be a square

$$CA = AP = PB = BC = 4$$
 cm

∴ Length of tangent = 4 cm.

11. : PQ is the tangent and OP is the radius through the point of contact.



$$\therefore \angle OPQ = 90^{\circ}$$

[The tangent at any point of a circle is perpendicular to the radius through the point of contact]

By Pythagoras theorem in right riangle OPQ ,

$$\mathrm{OQ}^2 = \mathrm{OP}^2 + \mathrm{PQ}^2$$

$$\Rightarrow$$
 $(12)^2 = (5)^2 + PQ^2$


$$\Rightarrow$$
 144 = 25 + PQ²

$$\Rightarrow PQ^2 = 144 - 25$$

$$\Rightarrow$$
 PQ² = 119

$$\Rightarrow PQ = \sqrt{119} \text{cm}$$

Hence, the length PQ is $\sqrt{119}$ cm.



In the given figure, PR and CR are both tangents drawn to the same circle from an external point R.

$$\therefore$$
 PR = CR.....(i)

Also, QR and CR are both tangents drawn to the same circle (second circle) from an external point R

$$QR = CR....(ii)$$

From (i) and (ii), we get

PR = QR [each equal to CR].

R is the midpoint of PQ,

i.e., the common tangent to the circles at C, bisects the common tangent at P and Q.

13. If
$$\angle PBO = 30^{\circ}$$

Then $\angle OPB = 30^o$ [angles opposite to equal sides are equal]

and $\angle OPT = 90^o$ [angle between radius and tangent]

$$\angle BPT = \angle OPT + \angle OPB$$

$$\angle BPT = 90^{\circ} + 30^{\circ} = 120^{\circ}$$


Now, $\angle PTA = 180^o - (\angle OBP + \angle BPT)$ [angle sum property of a triangle]

$$\angle PTA = 180^{o} - (30^{o} + 120^{o})$$

$$\angle PTA = 180^{\circ} - (150^{\circ})$$

$$\therefore \angle PTA = 30^{\circ}$$

14. In the given figure, O is the centre of a circle. PT and PQ are tangents to the circle from an external point P. If $\angle TPQ$ = 70°, then,we have to find $\angle TRQ$.



We know that the radius and tangent are perpendicular at their point of contact.

$$\angle OTP = \angle OQP = 90^{\circ}$$

Now, In quadrilateral OQPT

$$\angle QOT + \angle OTP + \angle OQP + \angle TPQ = 360^\circ$$
 [Angle sum property of a quadrilateral]


$$\angle QOT + 90 + 90 + 70 = 360$$

$$250 + \angle QOT = 360 =$$

$$\angle QOT = 110^{\circ}$$

We know that the angle subtended by an arc at the centre is double of the angle subtended by the arc at any point on the circumference of the circle.

$$\angle TRQ = \frac{1}{2} \angle QOT \Rightarrow \angle TRQ = \frac{1}{2} \times 110 = 55$$



Point P is the midpoint of arc \widehat{QR} of a circle with centre 0.

ST is the tangent to the circle at point P.

TO prove :Chord QR||ST

Proof: P is the midpoint of \widehat{QR}

$$\Rightarrow \widehat{QP} = \widehat{PR}$$


 \Rightarrow chord QP = chord PR [: in a circle, if two arcs are equal, then their corresponding chords are equal]

$$\therefore \angle PQR = \angle PRQ$$

$$\Rightarrow$$
 $\angle TPR = \angle PRQ$ [as , $\angle PQR = \angle TPR$, angles in alternate segments]

$$\Rightarrow$$
 $QR\|ST$, $[\because \angle TPR \text{ and } \angle PRQ \text{ are alternate interior } angles]$

16. Let 0 be the common centre of the two concentric circles.



Let AB be a chord of the larger circle which touches the smaller circle at P. Join OP and OA

Then, \angle OPA = 90° [: The tangent at any point of a circle is perpendicular to th radius through the point of contact]

$$\therefore$$
 OA² = OP² + AP² By Pythagoras theorem

$$\Rightarrow$$
 (5)² = (3)² + AP²

$$\Rightarrow$$
 25 = 9 + AP²

$$\Rightarrow$$
 P² = 25 - 9

$$\Rightarrow AP^2 = 16$$


$$\Rightarrow$$
 AP = $\sqrt{16}$ = 4 cm


Since the perpendicular from the centre of a circle to a chord bisects the chord, therfore,

$$AP = BP = 4 \text{ cm}$$

$$\therefore$$
 AB = AP + BP = AP + AP = 2AP = 2(4) = 8 cm

Hence, the required length is 8 cm.





TO PROVE
$$\angle AOB + \angle COD$$
= 180° and, $\angle AOD + \angle BOC$ = 180°

CONSTRUCTION Join OP, OQ, OR and OS.


PROOF Since the two tangents drawn from an external point to a circle subtend equal angles at the centre.

Sum of all the angles subtended at a point is 360°

$$\Rightarrow \quad 2(\angle 2 + \angle 3 + \angle 6 + \angle 7) = 360^{\circ} \text{ and } 2(\angle 1 + \angle 8 + \angle 4 + \angle 5) = 360^{\circ}$$

$$\Rightarrow \quad (\angle 2 + \angle 3) + (\angle 6 + \angle 7) = 180^{\circ} \text{ and } (\angle 1 + \angle 8) + (\angle 4 + \angle 5) = 180^{\circ}$$

$$\Rightarrow \quad \angle AOB + \angle COD = 180^{\circ} \begin{bmatrix} \because \angle 2 + \angle 3 = \angle AOB, \angle 6 + \angle 7 = \angle COD \\ \angle 1 + \angle 8 = \angle AOD \text{ and } \angle 4 + \angle 5 = \angle BOC \end{bmatrix}$$
and, $\angle AOD + \angle BOC = 180^{\circ}$



Let the sides BC, CA, AB of \triangle ABC touch the incircle at D, E, F respectively.

Join the centre O of the circle with A, B, C, D, E, F

Since, tangents to a circle from an external point are equal

$$\therefore$$
 CE = CD = 6 cm

$$BF = BD = 8 cm$$

$$AE = AF = x cm (say)$$

$$AB = (x + 8)$$
 cm and $AC = (x + 6)$ cm and $CB = 6 + 8 = 14$ cm

Area of
$$\triangle$$
 OAB = $\frac{1}{2}$ (8 + x) × 4 = (16 + 2x) cm²(i)

area of
$$\triangle$$
 OBC = $\frac{1}{2} \times 14 \times 4 = 28 \text{ cm}^2$(ii)

area of
$$\triangle$$
 OCA = $\frac{1}{2}$ (6 + x) × 4 = (12 + 2x) cm²(iii)

: area of
$$\triangle$$
 ABC = 16 + 2x + 12 + 2x + 28 = (4x + 56) cm² (iv)

Again, perimeter of $\triangle ABC = AC + AB + BC$

$$= 6 + x + (8 + x) + (6 + 8)$$

$$= 28 + 2x = 2(14 + x) \text{ cm}$$

$$S = \frac{2(14+x)}{2} = 14 + x$$

Area of
$$\triangle$$
 ABC = $\sqrt{s(s-a)(s-b)(s-c)}$

Area of
$$\triangle$$
 ABC = $\sqrt{s(s-a)(s-b)(s-c)}$
= $\sqrt{(14+x)(14+x-14)(14+x-6-x)(14+x-8-x)}$
= $\sqrt{(14+x)48x}$

$$\sqrt{672x + 48x^2}.....(v)$$

$$\therefore (4x + 56) = \sqrt{672x + 48x^2} [By 4 \text{ and } 5]$$

$$\Rightarrow (4x + 56)^2 = 672x + 48x^2$$

$$\Rightarrow 16(x + 14)^2 = 16(42x + 3x^2)$$

$$\Rightarrow (x + 14)^2 = (42x + 3x^2)$$

$$\Rightarrow x^2 + 28x + 196 = 3x^2 + 42x$$

$$(x + 14)(x - 7) = 0$$

$$x = 7, x = -14$$
But $x = -14$ is not possible.

But x = -14 is not possible

$$... x = 7$$

$$AB = x + 8 = 7 + 8 = 15 \text{ cm}$$

and
$$AC = x + 6 = 7 + 6 = 13$$
 cm


19. Given: ABCD is a quadrilateral circumscribing a circle whose centre is 0.

To prove:

i.
$$\angle AOB + \angle COD = 180^{\circ}$$

ii.
$$\angle BOC + \angle AOD = 180^{\circ}$$

Construction: Join OP, OQ, OR and OS.



Proof: Since tangents from an external point to a circle are equal.

$$\therefore$$
 AP = AS,

$$BP = BQ.....(i)$$

$$CQ = CR$$

$$DR = DS$$

In \triangle OBP and \triangle OBQ,

OP = OQ [Radii of the same circle]

OB = OB [Common]

BP = BQ [From eq. (i)]

 $\therefore \triangle$ OPB $\cong \triangle$ OBQ [By SSS congruence criterion]

$$\therefore$$
 $\angle 1 = \angle 2$ [By C.P.C.T.]

Similarly,
$$\angle 3 = \angle 4$$
, $\angle 5 = \angle 6$, $\angle 7 = \angle 8$

Since, the sum of all the angles round a point is equal to 360° .

$$\therefore \angle 1 + \angle 2 + \angle 3 + \angle 4 + \angle 5 + \angle 6 + \angle 7 + \angle 8 = 360^{\circ}$$

$$\Rightarrow$$
 $\angle 1 + \angle 1 + \angle 4 + \angle 4 + \angle 5 + \angle 5 + \angle 8 + \angle 8 = 360^{\circ}$

$$\Rightarrow$$
 2 ($\angle 1 + \angle 4 + \angle 5 + \angle 8$) = 360°


$$\Rightarrow \angle 1 + \angle 4 + \angle 5 + \angle 8 = 180^{\circ}$$


$$\Rightarrow$$
 ($\angle 1 + \angle 5$) + ($\angle 4 + \angle 8$) = 180°

$$\Rightarrow \angle AOB + \angle COD = 180^{\circ}$$

Similarly we can prove that

$$\angle BOC + \angle AOD = 180^{\circ}$$





So,
$$\angle OPO' = 90^{\circ}$$

In right angled \triangle OPO'

$$OP = 4 cm$$

$$O'P = 3 cm$$

By pythagoras theorem, we get

$$OO'^2 = OP^2 + O'P^2$$

$$=4^2+3^2$$

$$=16+9=25$$

$$OO' = 5cm$$
.

Let
$$O'T = x$$
, then $OT = 5 - x$

In right angled \triangle PTO

By pythagoras theorem, we get

$$OP^2 = OT^2 + PT^2$$

$$\Rightarrow PT^2 = OP^2 - OT^2 \ PT^2 = 4^2 - (5 - x)^2$$
...(i)

In right angled \triangle PTO'

By pythagoras theorem, we get

$$O'P^2 = O'T^2 + PT^2$$

$$\Rightarrow PT^2 = O'P^2 - O'T^2$$

$$PT^2 = 3^2 - x^2$$
...(ii)

From (i) and (ii), we get

$$3^2 - x^2 = 4^2 - (5 - x)^2$$

$$9 - x^2 = 16 - 25 - x^2 + 10x$$

$$18 = 10x$$

$$\Rightarrow x = \frac{18}{10} = 1.8$$

Substitute x in (ii), we get

$$PT^2 = 3^2 - 1.8^2 = 9 - 3.24 = 5.76$$

$$PT = \sqrt{5.76} = 2.4$$

$$\Rightarrow \mathrm{PQ} = 2\mathrm{PT}$$

$$=2\times2.4$$

$$\therefore PQ = 4.8cm.$$

